

Vitamin E-Acetate SD 50



Chemical names of active ingredient

DL- α -tocopheryl acetate,
DL-alpha-tocopherol acetate, all-rac-
alpha-tocopherol acetic acid ester, racemic
5,7,8-trimethyltolcol acetate

CAS-No. 52225-20-4

EINECS-No. 231-710-0

PRD-No.
30041058*

* The product is kosher.

Article

50047396 25 kg bag in box

Country of origin

Germany

Units

1 International Unit (IU) = 1 mg vitamin E-acetate

Description


White to creamy-white powder with a
characteristic mild odor.

Composition

Ingredients in descending order of weight:
vitamin E-acetate, lactose, sodium caseinate,
glycerin monostearate, tricalcium phosphate.

Solubility

The product can be dissolved both in cold (10 °C)
and warm (35 – 40 °C) water, to form a stable milky
emulsion. It is virtually insoluble in organic solvents.



$C_{31}H_{52}O_3$

Molar mass 472.8 g/mol

Specification

Assay min. 50% DL- α -tocopheryl acetate
(= 500 former IU of vitamin E =
336 α -TE per gram)

For further information see separate document:
"Standard Specification" (not for regulatory
purposes) available via BASF's WorldAccount:
<https://worldaccount.basf.com> (registered access).

Monographs

The active ingredient complies with the current
"All-rac- α -tocopheryl acetate" Ph. Eur., "Vitamin E"
USP, and "All-rac-alpha-tocopheryl acetate" FCC
monographs.

Regulations

The product meets the regulatory requirements
for a vitamin E source in most countries. How-
ever, regulations on the ingredients used in the
respective countries and for the intended use
have to be observed.

Bulk density

0.4 – 0.55 g/ml

Stability

Stored in its unopened original packaging at room
temperature (max. 25 °C), the product is stable
for at least 36 months.

Storage/Handling

The product should be stored tightly sealed in the original packaging at max. 25 °C, in a dry place.

Applications

Dietary supplements:

The product is suitable for the direct compression of chewable tablets, sugar- and film-coated tablets, as well as hard gelatin capsules. Due to its excellent dispersibility in cold water, it is also ideal for use in preparations, which need to be dissolved or dispersed in water prior to use, such as low-acid or acid-free instant-drink granules.

Food products:

Food products, e.g., childrens food, except for non-cereal-based Infant Food formulations in the EU, dietetic products, food products for athletes, dried milk, milk shake and cocoa powder, as well as vitamin mixtures for the food industry. The content of milk protein in the product may trigger a reaction with acid resulting in undesirable precipitates. Therefore, it is not suitable for the production of acidic juices, effervescent tablets, and instant-drink granules.

Note

Vitamin E-Acetate SD 50 must be handled in accordance with the Safety Data Sheet.

This document, or any answers or information provided herein by BASF, does not constitute a legally binding obligation of BASF. While the descriptions, designs, data and information contained herein are presented in good faith and believed to be accurate, it is provided for your guidance only. Because many factors may affect processing or application/use, we recommend that you make tests to determine the suitability of a product for your particular purpose prior to use. It does not relieve our customers from the obligation to perform a full inspection of the products upon delivery or any other obligation.

NO WARRANTIES OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH, OR THAT THE PRODUCTS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS. IN NO CASE SHALL THE DESCRIPTIONS, INFORMATION, DATA OR DESIGNS PROVIDED BE CONSIDERED A PART OF OUR TERMS AND CONDITIONS OF SALE.

November 2009